

EXTERIOR POWERS OF THE REFLECTION REPRESENTATION IN THE COHOMOLOGY OF SPRINGER FIBRES

ANTHONY HENDERSON

ABSTRACT. Let $H^*(\mathcal{B}_e)$ be the cohomology of the Springer fibre for the nilpotent element e in a simple Lie algebra $\mathfrak{g}$, on which the Weyl group W acts by the Springer representation. Let $\Lambda^i V$ denote the i th exterior power of the reflection representation of W . We determine the degrees in which $\Lambda^i V$ occurs in the graded representation $H^*(\mathcal{B}_e)$, under the assumption that e is regular in a Levi subalgebra and satisfies a certain extra condition which holds automatically if $\mathfrak{g}$ is of type A, B, or C. This partially verifies a conjecture of Lehrer–Shoji, and extends the results of Solomon in the $e = 0$ case and Lehrer–Shoji in the $i = 1$ case. The proof proceeds by showing that $(H^*(\mathcal{B}_e) \otimes \Lambda^* V)^W$ is a free exterior algebra on its subspace $(H^*(\mathcal{B}_e) \otimes V)^W$.

1. INTRODUCTION

Let $\mathfrak{g}$ be a simple complex Lie algebra of rank ℓ . Let W denote the Weyl group of $\mathfrak{g}$, and let V be the reflection representation of W . It is well known that the exterior powers $\Lambda^i V$, for $i = 0, 1, \dots, \ell$, are inequivalent irreducible representations of W , each of which is self-dual.

Let e be a nilpotent element of $\mathfrak{g}$. The Springer fibre $\mathcal{B}_e$ is the variety of Borel subalgebras of $\mathfrak{g}$ containing e . Let $H^*(\mathcal{B}_e)$ denote the graded cohomology ring of $\mathcal{B}_e$ with complex coefficients; the cohomology lives solely in even degrees, so $H^*(\mathcal{B}_e)$ is commutative. We have the Springer representation of W on each $H^{2j}(\mathcal{B}_e)$. See [Sho], [Hum, Chapter 9].

Let s (depending on e) denote the multiplicity of the irreducible representation V in the total representation $H^*(\mathcal{B}_e)$. Let $m_1, m_2, \dots, m_s$ be the multiset of nonnegative integers, listed in increasing order, which are the halved degrees of the occurrences of V in the graded representation $H^*(\mathcal{B}_e)$. That is, we have by definition

$$(1.1) \quad \sum_j \dim(H^{2j}(\mathcal{B}_e) \otimes V)^W q^j = q^{m_1} + q^{m_2} + \dots + q^{m_s},$$

The author's research is supported by ARC grant DP0985184.

an equality of polynomials in the indeterminate q .

In the special case $e = 0$, it is well known that $H^*(\mathcal{B})$ is isomorphic to the coinvariant algebra $C^*(W)$ of W , that $s = \ell$, and that $m_1, \dots, m_\ell$ are the exponents of W . More generally, if e is a regular nilpotent in a Levi subalgebra of semisimple rank r , then it was proved by Lehrer and Shoji in [LS, Theorem 2.4] (see also [Spa]) that $s = \ell - r$ and that $m_1, \dots, m_s$ are the coexponents of the corresponding parabolic hyperplane arrangement, in the sense of Orlik and Solomon. See [ST] for some other related interpretations of these coexponents. For $\mathfrak{g}$ of classical type and general e , the numbers $m_1, m_2, \dots, m_s$ were calculated by Spaltenstein in [Spa, Propositions 1.6–1.9].

Lehrer and Shoji conjectured that, at least in the parabolic case which they considered, the occurrences of each exterior power $\Lambda^i V$ in $H^*(\mathcal{B}_e)$ were also controlled in a natural way by $m_1, m_2, \dots, m_s$.

Conjecture 1.1. [LS, Conjecture 8.3] *Suppose that e is a regular nilpotent in a Levi subalgebra. Then for any $i = 0, 1, \dots, \ell$,*

$$\sum_j \dim(H^{2j}(\mathcal{B}_e) \otimes \Lambda^i V)^W q^j = e_i(q^{m_1}, q^{m_2}, \dots, q^{m_s}),$$

where the right-hand side means the i th elementary symmetric polynomial in $q^{m_1}, q^{m_2}, \dots, q^{m_s}$, which is defined to be zero if $i > s$.

The $e = 0$ case of this conjecture had already been proved by Solomon in [Sol]; indeed, he proved the stronger statement that the algebra $(C^*(W) \otimes \Lambda^* V)^W$ is a free exterior algebra on $(C^*(W) \otimes V)^W$.

In Section 2, we prove the following generalization of Solomon's result, which implies various cases of Conjecture 1.1.

Theorem 1.2. *Suppose that e is regular in a Levi subalgebra of $\mathfrak{g}$, and define s and $m_1, \dots, m_s$ as above. Also suppose that there is a parabolic subgroup W_K of W such that the following two conditions hold.*

- (1) *There exist invariant polynomials $f_1, f_2, \dots, f_s \in (S^* V)^W$, homogeneous of degrees $m_1 + 1, m_2 + 1, \dots, m_s + 1$, whose restrictions to the reflection representation V_K of W_K form a set of fundamental invariants for W_K .*
- (2) *The nilpotent orbit of e intersects the nilradical of the parabolic subalgebra $\mathfrak{p}_K$ associated to W_K .*

(See Section 2 for the definitions of V_K and $\mathfrak{p}_K$.) Then the algebra $(H^*(\mathcal{B}_e) \otimes \Lambda^* V)^W$ is a free exterior algebra on $(H^*(\mathcal{B}_e) \otimes V)^W$. More precisely, the natural algebra homomorphism

$$\psi : \Lambda^*((H^*(\mathcal{B}_e) \otimes V)^W) \rightarrow (H^*(\mathcal{B}_e) \otimes \Lambda^* V)^W$$

is an isomorphism.

Here the domain and codomain of ψ are $(\mathbb{N} \times \mathbb{N})$ -graded algebras over $\mathbb{C}$, where the (i, j) -components are

$$\Lambda^i((H^{2j}(\mathcal{B}_e) \otimes V)^W) \text{ and } (H^{2j}(\mathcal{B}_e) \otimes \Lambda^i V)^W$$

respectively, and in both cases the algebra multiplication is graded-commutative with respect to the $\mathbb{N}$ -grading labelled by i ; the homomorphism ψ is induced by the inclusion of the subspace $(H^*(\mathcal{B}_e) \otimes V)^W$ in $(H^*(\mathcal{B}_e) \otimes \Lambda^* V)^W$. Since the graded degrees of this subspace are $(1, m_1), (1, m_2), \dots, (1, m_s)$, the statement that ψ is an isomorphism implies that

$$\sum_{i,j} \dim(H^{2j}(\mathcal{B}_e) \otimes \Lambda^i V)^W t^i q^j = (1 + tq^{m_1})(1 + tq^{m_2}) \cdots (1 + tq^{m_s}),$$

which is equivalent to Conjecture 1.1.

To illustrate the scope of the assumptions in Theorem 1.2, we prove in Section 3 the following results about types A–C.

Proposition 1.3. *If $\mathfrak{g}$ is of type A, then the assumptions of Theorem 1.2 hold for any e .*

Proposition 1.4. *If $\mathfrak{g}$ is of type B, then the assumptions of Theorem 1.2 hold for any e which is regular in a Levi subalgebra.*

Proposition 1.5. *If $\mathfrak{g}$ is of type C, then the assumptions of Theorem 1.2 hold for any e which is regular in a Levi subalgebra.*

Hence Conjecture 1.1 is proved in types A–C. By contrast, suppose that $\mathfrak{g}$ is of type D_4 and e has Jordan type $(3^2 1^2)$ in the natural representation on $\mathbb{C}^8$. Then e is regular in a Levi subalgebra of type A_2 , but we have $m_2 = 2$ and there are no W -invariant polynomials of degree 3, so condition (1) of Theorem 1.2 cannot be satisfied.

Acknowledgements and historical comments. The main result of this paper, Theorem 1.2, dates from 1997, when I was a student at the University of Sydney, supervised by Gus Lehrer. As the reader will observe, it is indebted to Lehrer’s ideas, and I thank him for his help and encouragement. I did not publish this result at the time, since it did not prove the motivating Conjecture 1.1 in general. Recently Eric Sommers has completed the proof of Conjecture 1.1 by a different method, and also removed the assumption that e is regular in a Levi subalgebra. I thank him for his interest in my old result, and for the suggestion that it be published.

2. PROOF OF THEOREM 1.2

Continue the notation of the introduction. Let $\mathfrak{h} \subset \mathfrak{b}$ be a Cartan subalgebra and Borel subalgebra of $\mathfrak{g}$, and let $\Pi \subset \Phi^+ \subset \Phi$ be the corresponding set of simple roots, positive roots, and roots. We identify W with the subgroup of $GL(\mathfrak{h})$ generated by the simple reflections s_α for $\alpha \in \Pi$; the reflection representation V of W is merely $\mathfrak{h}$ itself.

Let $J \subseteq \Pi$ be a subset of size r , and set $s = \ell - r$. We have a Levi subalgebra $\mathfrak{l}_J$ and parabolic subalgebra $\mathfrak{p}_J$ containing $\mathfrak{h}$ and $\mathfrak{b}$ respectively, a parabolic subsystem Φ_J of Φ , and a parabolic subgroup W_J of W . Define

$$V^J = \bigcap_{\alpha \in J} \ker(\alpha) = V^{W_J}.$$

We write V_J for the unique W_J -invariant complement to V^J in V , which is the reflection representation of W_J . Note that $\dim V^J = s$ and $\dim V_J = r$. Let $\mathcal{A}^J$ and $\mathcal{A}_J$ be the hyperplane arrangements in V^J and V_J respectively induced by the root hyperplanes in V .

We assume for the remainder of the section that

$$(2.1) \quad e \text{ is parabolic of type } J,$$

meaning that the orbit of e contains the regular nilpotent elements of $\mathfrak{l}_J$. As mentioned in the introduction, Lehrer and Shoji proved in this case that

$$(2.2) \quad \sum_j \dim(H^{2j}(\mathcal{B}_e) \otimes V)^W q^j = q^{m_1} + q^{m_2} + \cdots + q^{m_s},$$

where $m_1, \dots, m_s$ are the coexponents of the arrangement $\mathcal{A}^J$. (See [LS, Theorem 2.4]; the missing case in type D is covered by the results of Spaltenstein [Spa].)

An important special feature of the parabolic case is Lusztig's Induction Theorem for Springer representations.

Theorem 2.1. [Lus] *Under the assumption (2.1), the representation of W on $H^*(\mathcal{B}_e)$, neglecting the grading, is isomorphic to the induction $\text{Ind}_{W_J}^W(\mathbb{C})$ of the trivial representation of W_J .*

Corollary 2.2. *Under the assumption (2.1), we have*

$$\dim(H^*(\mathcal{B}_e) \otimes \Lambda^* V)^W = 2^s,$$

so the domain and codomain of ψ have the same dimension.

Proof. By Frobenius Reciprocity, we know that

$$(2.3) \quad \dim(\text{Ind}_{W_J}^W(\mathbb{C}) \otimes \Lambda^* V)^W = \dim(\Lambda^* V)^{W_J}.$$

From the fact that $(\Lambda^* V)^{s_\alpha} = \Lambda^*(\ker(\alpha))$ for all $\alpha \in J$ one deduces $(\Lambda^* V)^{W_J} = \Lambda^*(V^J)$, and the Corollary follows. $\square$

So to prove Theorem 1.2, it suffices to show that the homomorphism ψ is injective. Since the domain is an exterior algebra, this will follow if we can show that

$$(2.4) \quad \psi(\Lambda^s((H^*(\mathcal{B}_e) \otimes V)^W)) \neq 0.$$

To be more concrete, let $v_1, \dots, v_\ell$ be a basis of V , and let $c_{i,j} \in H^{2m_i}(\mathcal{B}_e)$, for $1 \leq i \leq s$, $1 \leq j \leq \ell$, be such that

$$\sum_{j=1}^n c_{1,j} \otimes v_j, \sum_{j=1}^n c_{2,j} \otimes v_j, \dots, \sum_{j=1}^n c_{s,j} \otimes v_j$$

is a basis of $(H^*(\mathcal{B}_e) \otimes V)^W$. Then by definition of ψ , (2.4) is equivalent to the statement that for some sequence $1 \leq j_1 < j_2 < \dots < j_s \leq n$,

$$(2.5) \quad \det \begin{pmatrix} c_{1,j_1} & c_{1,j_2} & \cdots & c_{1,j_s} \\ c_{2,j_1} & c_{2,j_2} & \cdots & c_{2,j_s} \\ \vdots & \vdots & \ddots & \vdots \\ c_{s,j_1} & c_{s,j_2} & \cdots & c_{s,j_s} \end{pmatrix} \neq 0.$$

To handle such a determinant, we need a better grasp of the cohomology ring $H^*(\mathcal{B}_e)$ in which it lives.

Using the W -equivariant isomorphism of V with its dual, we will identify the symmetric algebra S^*V with the ring of polynomial functions on V . It is well known that the invariant subring $(S^*V)^W$ is freely generated by ℓ homogeneous polynomials, called fundamental invariants for W . The coinvariant algebra $C^*(W)$ of W is the quotient S^*V/I , where I is the ideal generated by these fundamental invariants.

Now there is a canonical (degree-doubling) W -equivariant algebra homomorphism $S^*V \rightarrow H^*(\mathcal{B})$ which identifies $H^*(\mathcal{B})$ with $C^*(W)$ (see [Sho, §5]). Composing this with the natural homomorphism $H^*(\mathcal{B}) \rightarrow H^*(\mathcal{B}_e)$, we obtain a W -equivariant homomorphism $\varphi : S^*V \rightarrow H^*(\mathcal{B}_e)$ (see [HS, Theorem 1.1]). Note that the image of φ is contained in the subspace $H^*(\mathcal{B}_e)^{A(e)}$ of invariants for the component group of the centralizer of e in the adjoint group of $\mathfrak{g}$; in particular, φ is not surjective in general. However, it may happen that the induced map $(S^*V \otimes V)^W \rightarrow (H^*(\mathcal{B}_e) \otimes V)^W$ is surjective even if φ is not. We will see that this occurs under the assumptions of Theorem 1.2, which means that a calculation with polynomials on V suffices to prove (2.5).

Henceforth we let $K \subseteq \Pi$ be a subset satisfying conditions (1) and (2) of Theorem 1.2. Note that condition (1) entails $|K| = s$. Choose a basis $v_1, v_2, \dots, v_\ell$ of V such that $v_1, \dots, v_s$ is a basis of V_K . Since the

exterior derivative $S^*V \rightarrow S^*V \otimes V : f \mapsto \sum \frac{\partial f}{\partial v_j} \otimes v_j$ is W -equivariant, we have the following s elements of $(H^*(\mathcal{B}_e) \otimes V)^W$:

$$\sum_{j=1}^n \varphi\left(\frac{\partial f_1}{\partial v_j}\right) \otimes v_j, \sum_{j=1}^n \varphi\left(\frac{\partial f_2}{\partial v_j}\right) \otimes v_j, \dots, \sum_{j=1}^n \varphi\left(\frac{\partial f_s}{\partial v_j}\right) \otimes v_j.$$

We can prove simultaneously that these form a basis of $(H^*(\mathcal{B}_e) \otimes V)^W$, and that (2.5) holds (with $j_i = i$), by proving the single fact that $\varphi(\Delta) \neq 0$, where

$$\Delta = \det \begin{pmatrix} \frac{\partial f_1}{\partial v_1} & \frac{\partial f_1}{\partial v_2} & \dots & \frac{\partial f_1}{\partial v_s} \\ \frac{\partial f_2}{\partial v_1} & \frac{\partial f_2}{\partial v_2} & \dots & \frac{\partial f_2}{\partial v_s} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_s}{\partial v_1} & \frac{\partial f_s}{\partial v_2} & \dots & \frac{\partial f_s}{\partial v_s} \end{pmatrix}.$$

Since $v_1, \dots, v_s$ span the reflection representation of W_K , we have $w\Delta = \varepsilon(w)\Delta$ for all $w \in W_K$. This forces Δ to be divisible by the polynomial $\pi_K = \prod_{\beta \in \Phi_K^+} \beta$. Condition (1) tells us that on restricting to V_K , Δ becomes the Jacobian matrix of the fundamental invariants of W_K , which is well known to be a nonzero scalar multiple of the restriction to V_K of π_K (see [Ste]). So Δ is a nonzero scalar multiple of π_K , and it suffices to prove that $\varphi(\pi_K) \neq 0$.

By condition (2), we may suppose that e lies in the nilradical of $\mathfrak{p}_K$. Then any Borel subalgebra contained in $\mathfrak{p}_K$ must contain e , so we have an inclusion $\mathcal{B}^K \hookrightarrow \mathcal{B}_e$, where $\mathcal{B}^K$ denotes the variety of Borel subalgebras contained in $\mathfrak{p}_K$, which can be identified with the flag variety of $\mathfrak{l}_K$. Hence it suffices to prove that π_K is not in the kernel of the composition $S^*V \rightarrow H^*(\mathcal{B}) \rightarrow H^*(\mathcal{B}^K)$. But this composition is the canonical homomorphism identifying $C^*(W_K)$ with $H^*(\mathcal{B}^K)$, which maps π_K to a generator of the top-degree cohomology of $\mathcal{B}^K$ (see [HS, Proposition 1.4], which uses exactly this argument in the case when e lies in the Richardson orbit of $\mathfrak{p}_K$). This completes the proof of Theorem 1.2.

3. PROOFS OF PROPOSITIONS 1.3–1.5

Proof of Proposition 1.3. In type A, we have $\mathfrak{g} = \mathfrak{sl}_{\ell+1}$ and $W = S_{\ell+1}$. The simple roots are given by $\alpha_i = x_i - x_{i+1}$, where $x_1, \dots, x_{\ell+1}$ denote coordinate functions on V which are permuted by W , and which satisfy $x_1 + \dots + x_{\ell+1} = 0$. The orbit of the nilpotent e is determined by its Jordan type λ , which can be an arbitrary partition of $\ell+1$: the element e is then automatically parabolic of type

$$J = \Pi \setminus \{\alpha_{\lambda_1 + \lambda_2 + \dots + \lambda_i} \mid 1 \leq i < \ell(\lambda)\}, \text{ so } r = \ell - \ell(\lambda) + 1, s = \ell(\lambda) - 1.$$

In this case, $\mathcal{A}^J$ is an arrangement of type A_s , so $m_i = i$ for $i = 1, \dots, s$. We define f_i to be the usual $S_{\ell+1}$ -invariant polynomial of degree $i+1$, namely the $(i+1)$ th elementary symmetric polynomial in $x_1, \dots, x_{\ell+1}$. Let

$$K = \{\alpha_{r+1}, \alpha_{r+2}, \dots, \alpha_\ell\}.$$

Then V_K is defined by the equations $x_i = 0$ for all $i \leq r$, so the restrictions of the f_i 's to V_K are the elementary symmetric polynomials in $x_{r+1}, \dots, x_{\ell+1}$, and condition (1) is satisfied. With this K , condition (2) amounts to saying that there is a partial flag in the natural representation,

$$0 = U_0 \subset U_1 \subset U_2 \subset \dots \subset U_r \subset \mathbb{C}^{\ell+1},$$

such that $eU_i \subseteq U_{i-1}$ and $\dim U_i = i$ for all $i \geq 1$, and $e(\mathbb{C}^{\ell+1}) \subseteq U_r$. This is true: we can take $U_r = e(\mathbb{C}^{\ell+1})$, and then let $U_1 \subset \dots \subset U_r$ be any e -stable complete flag in U_r . $\square$

Proof of Proposition 1.4. In type B, we have $\mathfrak{g} = \mathfrak{o}_{2\ell+1}$. The simple roots are given by $\alpha_i = x_i - x_{i+1}$ for $i \leq \ell - 1$ and $\alpha_\ell = x_\ell$, where $x_1, \dots, x_\ell$ denote coordinate functions on V which are permuted up to sign by W . We are assuming that e is parabolic of type J , for some subset $J \subseteq \Pi$. There is some nonnegative integer $m \leq \ell$ and composition λ of $\ell - m$ such that

$$J = \Pi \setminus \{\alpha_{\lambda_1 + \lambda_2 + \dots + \lambda_i} \mid 1 \leq i \leq \ell(\lambda)\}, \text{ so } r = \ell - \ell(\lambda), s = \ell(\lambda).$$

The orbit containing the regular nilpotents in $\mathfrak{l}_J$ consists of those whose Jordan type in the natural representation $\mathbb{C}^{2\ell+1}$ is the partition obtained by rearranging $\lambda_1, \lambda_1, \lambda_2, \lambda_2, \dots, \lambda_s, \lambda_s, 2m+1$. Grouping the Jordan blocks together as appropriate, we get a direct sum decomposition $\mathbb{C}^{2\ell+1} = U \oplus U' \oplus U''$ where U and U' are $(\ell - m)$ -dimensional e -stable isotropic subspaces on each of which the Jordan type of e is the partition obtained by rearranging λ , U'' is a $(2m+1)$ -dimensional e -stable subspace on which e has a single Jordan block, and U'' is perpendicular to $U \oplus U'$ for the nondegenerate symmetric form.

The arrangement $\mathcal{A}^J$ is of type B_s , so $m_i = 2i-1$ for $i = 1, \dots, s$. We define f_i to be the usual W -invariant polynomial of degree $2i$, namely the i th elementary symmetric polynomial in $x_1^2, \dots, x_\ell^2$. Let

$$K = \{\alpha_{r+1}, \alpha_{r+2}, \dots, \alpha_\ell\}.$$

Then V_K is defined by the equations $x_i = 0$ for $i \leq r$, so the restrictions of the f_i 's to V_K are the elementary symmetric polynomials in $x_{r+1}^2, \dots, x_\ell^2$, and condition (1) is satisfied. With this K , condition (2)

amounts to saying that there is an isotropic partial flag in the natural representation,

$$0 = U_0 \subset U_1 \subset \cdots \subset U_r \subset \mathbb{C}^{2\ell+1},$$

such that $eU_i \subseteq U_{i-1}$ and $\dim U_i = i$ for all $i \geq 1$, and $e(U_r^\perp) \subseteq U_r$. This is true: if we define U_r to be $e(U) \oplus U'''$ where U''' is the unique e -stable m -dimensional subspace of U'' , then $U_r^\perp = U \oplus \ker(e|_{U'}) \oplus e^{-1}(U''')$, so $e(U_r^\perp) = U_r$. We can then let $U_1 \subset \cdots \subset U_r$ be any e -stable complete flag in U_r . $\square$

Proof of Proposition 1.5. In type C, we have $\mathfrak{g} = \mathfrak{sp}_{2\ell}$ with natural representation $\mathbb{C}^{2\ell}$; the description of the simple roots is the same as in type B, except that $\alpha_\ell = 2x_\ell$. The proof is identical to that of Proposition 1.4, but with the dimension of U'' (which equals the size of a Jordan block of e) changed from $2m+1$ to $2m$, so in the last step one has $U_r^\perp = U \oplus \ker(e|_{U'}) \oplus U'''$, still implying $e(U_r^\perp) \subseteq U_r$. $\square$

REFERENCES

- [HS] R. Hotta and T. A. Springer, *A specialization theorem for certain Weyl group representations and an application to the Green polynomials of unitary groups*, Invent. Math. **41** (1977), 113–127.
- [Hum] J. E. Humphreys, *Conjugacy Classes in Semisimple Algebraic Groups*, Mathematical Surveys and Monographs, Vol. 43, American Mathematical Society, Providence, Rhode Island (1995).
- [LS] G. I. Lehrer and T. Shoji, *On flag varieties, hyperplane complements and Springer representations of Weyl groups*, J. Austral. Math. Soc. (Series A) **49** (1990), no. 3, 449–485.
- [Lus] G. Lusztig, *An induction theorem for Springer’s representations*, in *Representation Theory of Algebraic Groups and Quantum Groups*, Adv. Stud. Pure Math. **40**, Math. Soc. Japan, Kinokuniya (2004), 253–259.
- [Sho] T. Shoji, *Geometry of orbits and Springer correspondence*, in *Orbites unipotentes et representations, I*, Astérisque **168**, Soc. Math. de France, Paris (1988), 61–140.
- [Sol] L. Solomon, *Invariants of finite reflection groups*, Nagoya Math. J. **22** (1963), 57–64.
- [ST] E. Sommers and P. Trapa, *The adjoint representation in rings of functions*, Represent. Theory **1** (1997), 182–189.
- [Spa] N. Spaltenstein, *On the reflection representation in Springer’s theory*, Comment. Math. Helvetici **66** (1991), no. 4, 618–636.
- [Ste] R. Steinberg, *Invariants of finite reflection groups*, Canad. J. Math. **12** (1960), 616–618.

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SYDNEY NSW
2006, AUSTRALIA

E-mail address: anthonyh@maths.usyd.edu.au